

Specimen Label

Renovate® OTF

On Target Flakes



Aquatic Sites: For control of emersed, submersed and floating aquatic plants in the following aquatic sites: ponds; lakes; reservoirs; marshes; wetlands; impounded rivers, streams and other bodies of water that are quiescent; non-irrigation canals, seasonal irrigation waters and ditches which have little or no continuous outflow.

Active Ingredient:

triclopyr: 3,5,6-trichloro-2-pyridinyloxyacetic acid,
triethylamine salt 14.0%
Other Ingredients 86.0%
TOTAL 100.0%

Acid equivalent: triclopyr - 10.0% - 1.6 ounces per pound.

Keep Out of Reach of Children

CAUTION/PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

ENVIRONMENTAL HAZARDS

Under certain conditions, treatment of aquatic weeds can result in oxygen depletion or loss due to decomposition of dead plants, which may contribute to fish suffocation. This loss can cause fish suffocation. Therefore, to minimize this hazard, do not treat more than one-third to one-half of the water area in a single operation and wait at least 10 to 14 days between treatments. Begin treatment along the shore and proceed outwards in bands to allow fish to move into untreated areas. Consult with the State agency for fish and game before applying to public water to determine if a permit is needed.

First Aid

If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 - 20 minutes.• Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.

EMERGENCY NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call **INFOTRAC** at **1-800-535-5053**.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Refer to label booklet for additional precautionary information and Directions for Use.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read "Warranty Disclaimer", "Inherent Risks of Use", and "Limitation of Remedies" at end of label booklet. If terms are unacceptable, return at once unopened.**

If you wish to obtain additional product information, please visit our web site at www.sepro.com.

EPA Reg. No. 67690-42
FPL 103006

Renovate is a registered trademark of Dow AgroSciences LLC.
Manufactured by: **SePRO Corporation** Carmel, IN 46032 U.S.A.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all *Directions for Use* carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

General Information

When applying this product follow all applicable use directions, precautions and limitations.

AQUATIC AND WETLAND SITES

Use Renovate® OTF herbicide for control of emerged, submersed and floating aquatic plants in the following aquatic sites: ponds; lakes; reservoirs; marshes; wetlands; impounded rivers, streams and other bodies of water that are quiescent; non-irrigation canals, seasonal irrigation waters and ditches which have little or no continuous outflow.

Obtain Required Permits: Consult with appropriate state or local water authorities before applying this product in and around public waters. State or local public agencies may require permits.

Recreational Use of Water in Treatment Area: There are no restrictions on use of water in the treatment area for recreational purposes, including swimming and fishing.

Livestock Use of Water from Treatment Area: There are no restrictions on livestock consumption of water from the treatment area.

GENERAL USE PRECAUTIONS AND RESTRICTIONS

Chemigation: Do not apply this product through any type of irrigation system.

Irrigation: Water treated with Renovate OTF may not be used for irrigation purposes for 120 days after application or until triclopyr residue levels are determined by laboratory analysis, or other appropriate means of analysis, to be 1.0 ppb or less.

Seasonal Irrigation Waters: Renovate OTF may be applied during the off-season to surface waters that are used for irrigation on a seasonal basis, provided that there is a minimum of 120 days between Renovate OTF application and the first use of treated water for irrigation purposes or until triclopyr residue levels are determined by laboratory analysis, or other appropriate means of analysis, to be 1.0 ppb or less.

Irrigation Canal/Ditches: DO NOT apply Renovate OTF to irrigation canals/ditches unless the 120 day restriction on irrigation water usage can be observed or triclopyr residue levels are determined by laboratory analysis, or other appropriate means of analysis, to be 1.0 ppb or less.

- **Do not** apply Renovate OTF directly to, or otherwise permit it to come into direct contact with grapes, tobacco, vegetable crops, flowers, or other desirable broadleaf plants, and do not permit granule dust to drift into them.
- **Do not** apply to salt water bays or estuaries.
- **Do not** apply directly to un-impounded rivers or streams.
- **Do not** apply on ditches or canals currently being used to transport irrigation water or that will be used for irrigation within 120 days following treatment or until triclopyr residue levels are determined to be 1.0 ppb or less.
- **Do not** apply where runoff water may flow onto agricultural land as injury to crops may result.

Grazing and Haying Restrictions:

Except for lactating dairy animals, there are no grazing restrictions following application of this product.

- **Grazing Lactating Dairy Animals:** **Do not** allow lactating dairy animals to graze treated areas until the next growing season following application of this product.
- **Do not** harvest hay for 14 days after application.
- Grazed areas of non-cropland and forestry sites may be spot treated if they comprise no more than 10% of the total grazable area.

Slaughter Restrictions: During the season of application, withdraw livestock from grazing treated grass at least 3 days before slaughter.

BEST MANAGEMENT PRACTICES FOR DRIFT MANAGEMENT

Equipment used in the application of Renovate OTF should be carefully calibrated before use and checked frequently during application to be sure it is working properly and delivering a uniform distribution pattern. To prevent increased Renovate OTF dosage above specified limits, do not overlap applications. Aerial application should be made only when the wind velocity is 2 to 10 mph.

Applications should be made only when there is little or no hazard for volatility or dust drift, and when application can maintain Renovate OTF placement in the intended area. Very small quantities of dust, which may not be visible, may seriously injure susceptible plants, and Renovate OTF may be blown outside of the intended treatment area under extreme conditions. Do not spread Renovate OTF when wind is blowing toward susceptible crops or ornamental plants that are near enough to be injured.

Avoiding drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for drift. The applicator is responsible for considering all these factors when making decisions.

Ground Application Equipment: To aid in reducing drift, Renovate OTF should be applied when wind velocity is low (follow state regulations; see Sensitive Area under *Aerial Drift Reduction Advisory* below).

AERIAL DRIFT REDUCTION ADVISORY

This section is advisory in nature and does not supersede the mandatory label requirements.

Application Height: Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces drift potential.

Swath Adjustment: When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (e.g. higher wind).

Wind: Drift potential is lowest between wind speeds of 2 - 10 mph (follow state regulations). However, many factors, including equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential.

Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

Sensitive Areas: Renovate OTF should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

AQUATIC WEEDS CONTROLLED BY RENOVATE OTF

- | | |
|-----------------------|---|
| alligatorweed | pennywort |
| American lotus | smartweed |
| bladderwort | water chestnut† |
| Eurasian watermilfoil | yellow water lily (<i>Nuphar spp.</i> , spatterdock) |
| milfoil species | white water lily (<i>Nymphaea spp.</i>) |
| parrotfeather†† | water primrose (<i>Ludwigia spp.</i>) |
| pickerelweed | watershield (<i>Brasenia spp.</i>) |

†Not for use in California.

††Retreatment may be needed to achieve desired level of control.

Application Methods

SURFACE APPLICATION

Use a mechanical spreader such as a fertilizer spreader or mechanical seeder or similar equipment capable of uniformly applying Renovate OTF. Before spreading any product, carefully calibrate the application equipment. When using boats and power equipment, you must determine the proper combination of (1) boat speed (2) rate of delivery from the spreader, and (3) width of swath covered by the granules.

Use the following formula to calibrate the spreader’s delivery in pounds of Renovate OTF per minute:

$$\frac{\text{Miles per hour} \times \text{spreader width (feet)} \times \text{X pounds per acre}}{495} = \text{Pounds per minute}$$

AERIAL APPLICATION (HELICOPTER ONLY)

Ensure uniform application. All equipment should be properly calibrated using blanks with similar physical characteristics to Renovate OTF. To avoid streaked, uneven or overlapped application, use an appropriate tracking device (e.g. GPS). Refer to the *Aerial Drift Reduction Advisory* section of this label for additional precautions and instructions for aerial application.

Floating and Emerged Weeds

For control of water lily’s (*Nymphaea spp.* and *Nuphar spp.*), watershield (*Brasenia spp.*), and other susceptible emerged and floating herbaceous weeds, apply 0.75 to 2.5 ppm triclopyr per acre. Apply when plants are actively growing.

Use higher rates in the rate range when plants are mature, when the weed mass is dense, or for difficult to control species. Repeat as necessary to control regrowth, but do not exceed a total of 2.5 ppm triclopyr for the treatment area per annual growing season.

Submersed Weeds

For control of Eurasian watermilfoil (*Myriophyllum spicatum*) and other susceptible submersed weeds

in ponds, lakes, reservoirs, impounded rivers, streams and other bodies of water that are quiescent; non-irrigation canals, and seasonal irrigation waters, or ditches that have little or no continuous outflow, apply Renovate OTF using mechanical or portable granule spreading equipment. Rates should be selected according to the rate chart below to provide a triclopyr concentration of 0.75 to 2.5 ppm ae in treated water. Use of higher rates in the rate range is recommended in areas of greater water exchange or when treating target area of 1/2 acre or smaller. These areas may require a repeat application. However, total application of Renovate OTF must not exceed an application rate of 2.5 ppm ae triclopyr for the treatment area per annual growing season.

For optimal control, apply in spring or early summer when Eurasian watermilfoil or other submersed weeds are actively growing.

Concentration of Triclopyr Acid in Water (ppm a.e.)					
Avg. Water Depth (ft)	Pounds Renovate OTF / acre (10% a.e.)				
	0.75 ppm	1.0 ppm	1.5 ppm	2.0 ppm	2.5 ppm
1	20	27	41	54	68
2	41	54	81	108	135
3	61	81	122	162	203
4 feet or greater	81	108	164	216	270

Precautions for Potable Water Intakes:

For applications of Renovate OTF to control floating, emerged, and submersed weeds in sites that contain a functioning potable water intake for human consumption, see the chart on the next page to determine the minimum setback distances of the application from the functioning potable water intakes.

Concentration of Triclopyr Acid in Water (ppm a.e.)

Area Treated (acres)	Required Setback Distance (ft) from Potable Water Intake				
	0.75 ppm	1.0 ppm	1.5 ppm	2.0 ppm	2.5 ppm
<4	300	400	600	800	1000
>4 - 8	420	560	840	1120	1400
>8 - 16	600	800	1200	1600	2000
>16 - 32	780	1040	1560	2080	2600
>32 acres, calculate a setback using the formula for the appropriate rate	Setback (ft) = $\frac{(800 \cdot \ln(\text{acres}) - 160)}{3.33}$	Setback (ft) = $\frac{(800 \cdot \ln(\text{acres}) - 160)}{2.50}$	Setback (ft) = $\frac{(800 \cdot \ln(\text{acres}) - 160)}{1.67}$	Setback (ft) = $\frac{(800 \cdot \ln(\text{acres}) - 160)}{1.25}$	Setback (ft) = $\frac{(800 \cdot \ln(\text{acres}) - 160)}{1.00}$

Note: ln = natural logarithm

Example Calculation 1:

to apply 2.5 ppm Renovate OTF to 50 acres:

$$\begin{aligned}\text{Setback in feet} &= (800 \times \ln(50 \text{ acres}) - 160) \\ &= (800 \times 3.912) - 160 \\ &= 2970 \text{ feet}\end{aligned}$$

Example Calculation 2:

to apply 0.75 ppm Renovate OTF to 50 acres:

$$\begin{aligned}\text{Setback in feet} &= \frac{(800 \times \ln(50 \text{ acres}) - 160)}{3.33} \\ &= \frac{(800 \times 3.912) - 160}{3.33} \\ &= 892 \text{ feet}\end{aligned}$$

Note: Existing potable water intakes which are no longer in use, such as those replaced by potable water wells or connections to a municipal water system, are not considered to be functioning potable water intakes.

To apply Renovate OTF around and within the distances noted above from a functioning potable water intake, the intake must be turned off until the triclopyr level in the intake water is determined to be 0.4 parts per million (ppm) or less by laboratory analysis or immunoassay.

WETLAND SITES

Wetlands include flood plains, deltas, marshes, swamps, bogs, and transitional areas between upland and lowland sites. Wetlands may occur within forests, wildlife habitat restoration and management areas and similar sites as well as areas adjacent to or surrounding domestic water supply reservoirs, lakes and ponds.

For control of emersed, floating or submersed aquatic weeds in wetland sites, follow use directions and application methods associated with the *Floating and Emersed Weeds* or *Submersed Weeds* sections on this label.

Use Precautions

Minimize unintentional application to open water when treating target vegetation in wetland sites. **Note:** Consult local public water control authorities before applying this product in and around public water. Permits may be required to treat such areas.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

SePRO Corporation warrants that the product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SEPRO CORPORATION MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of SePRO Corporation as the seller. To the extent permitted by applicable law all such risks shall be assumed by buyer.

Limitation of Remedies

To the fullest extent permitted by law, SePRO Corporation shall not be liable for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories) shall be limited to, at SePRO Corporation's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

SePRO Corporation shall not be liable for losses or damages resulting from handling or use of this product unless SePRO Corporation is promptly notified of such losses or damages in writing. In no case shall SePRO Corporation be liable for consequential or incidental damages or losses.

The terms of the "Warranty Disclaimer" above and this "Limitation of Remedies" cannot be varied by any written or verbal statements or agreements. No employee or sales agent of SePRO Corporation or the seller is authorized to vary or exceed the terms of the "Warranty Disclaimer" or "Limitations of Remedies" in any manner.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal. Open dumping is prohibited.

Pesticide Storage: Store in original container. Do not store near food or feed. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Disposal (Plastic Bags): Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Container Disposal (Plastic): Do not reuse container. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General: Consult federal, state, or local disposal authorities for approved alternative procedures.